

## CRINOIDEA CARE

SOP# = ECHI2


**PURPOSE:** To describe methods of care for feather stars.

**POLICY:** To provide optimum care for all animals.

**RESPONSIBILITY:** Collector and user of animals. If these are not the same person, user takes over responsibility of animals as soon as the animals have arrived on station.

**PROCEDURE:** At present there is one species of feather star found around BMSC.

### IDENTIFICATION:

Common Name	Scientific Name	Identifying Features
Feather Star	<i>Florometra serratissima</i>	<ul style="list-style-type: none"><li>- Ten long arms each with slender branches</li><li>- Color is tan to reddish tan</li><li>- Can reach 25 cm in height</li><li>- Found on soft and hard bottoms in shallow subtidal areas and to depths of 990m</li><li>- It is essentially an upside-down brittle star</li></ul> 

### CAPTURE:

Location	-Subtidal sites are found on the shores of the Deer Group Islands, Dixon I., Scott's Bay, off the Blowhole, along Grappler Inlet and at the Harbor mouth across from Aguilar Pt.
Methods	<ul style="list-style-type: none"><li>- Can be collected by hand from shallow sites and placed in a container</li><li>- From deeper sites must be collected by scuba diving</li></ul>
Important Notes	<b>** Be very careful when moving the feather stars, as their arms will break off very easily when roughly handled.</b>

## ANIMAL CARE:

<b>Holding Requirements</b>	<ul style="list-style-type: none"><li>- Held in continually flowing seawater</li><li>- Provide variety of rocks, as feather stars like to anchor to a roughened rock surface</li><li>- They will move around until they find a rock that they like. Lids are unnecessary once specimen is anchored</li></ul>
<b>Feeding</b>	<ul style="list-style-type: none"><li>- Feather stars are filter feeders</li><li>- They require supplemental feedings of plankton every 3<sup>rd</sup> to 5<sup>th</sup> day, as there is not enough plankton in BMSC's intake seawater</li></ul>
<b>Tank Cleaning</b>	<ul style="list-style-type: none"><li>- Once a month the feather stars should be removed from the tank and placed into a holding bucket</li><li>- <b>Be very careful when moving the feather stars, as their arms will break off very easily when roughly handled</b></li><li>- The tanks should be drained and the sides and bottom should be scrubbed first with warm freshwater</li><li>- Then rinsed with seawater and allowed to refill</li><li>- Place the feather stars back in the tank once refilling has begun.</li></ul>
<b>Daily Activities</b>	<ul style="list-style-type: none"><li>-Ensure water is flowing into the tank at a reasonable rate.</li><li>-Ensure the standpipe is in place and not blocked.</li><li>-Check for and remove dead animals.</li><li>-Check for and remove any uneaten prey organisms.</li><li>-Check for and remove foreign organisms.</li></ul>

SOP# = ECH12  
K. Bartlett  
2017